

A-Calm

Tablet

Tolperisone Hydrochloride

COMPOSITION

Each film-coated tablet contains Tolperisone Hydrochloride INN 50 mg.

PHARMACOLOGY

Tolperisone hydrochloride is a centrally acting muscle relaxant, which acts on the central nervous system and used mainly for the treatment of elevated muscle tone and tension as well as for certain circulatory problem in the extremities. It reduces experimental hypertonia and decerebration and is independent from the cerebral centers of vasoregulation.

INDICATIONS

Increased tone of skeletal muscles due to organic neurological disorders (e.g. injury of the pyramidal tract, multiple sclerosis, myelopathy, encephalomyelitis etc.), muscular hypertension, muscular spasm, muscular contraction & rigidity, spinal automatism and discopathy. Obliterate vascular diseases (obliterate arteriosclerosis, diabetic angiopathy, obliterative thromboangitis, raynaud's disease, diffusion scleroderma) and disorders due to injury innervation of the vessels (acrocyanosis, intermittent angioneurotic dysbasia). In individual cases post thrombotic venous and circulation disorders and crural ulcer.

DOSAGE & ADMINISTRATION

Route of administration: Oral.

One should follow the doctor's instructions regarding the dosage of Tolperisone hydrochloride. The recommended dose is as followings :

Adult : 50-150 mg three times daily according to the individual requirement and tolerance of the patients.

Children (from 3 months to 6 years) : 5-10 mg/kg/day in three divided doses.

OR AS DIRECTED BY THE PHYSICIAN.

CONTRAINDICATIONS

Hypersensitivity to tolperisone, Myasthenia gravis, nursing mother.

WARNING & PRECAUTIONS

Incase of children, the prescribed dose and duration of treatment should closely be observed.

SIDE-EFFECTS

Sometimes muscular weakness, headache, nausea, vomiting, and rarely hypersensitivity reactions.

USE IN PREGNANCY & LACTATION

Tolperisone Hydrochloride should be used in pregnancy to physician's advice. Tolperisone Hydrochloride should not be used during breast-feeding.

DRUG INTERACTIONS

Concomitant use of methocarbamol with Tolperisone Hydrochloride has been reported to cause disturbance of visual accommodation.

STORAGE

Store below 30° C temperature & dry place protected from light.
Keep all medicines out of reach of children.

PACKING

Each box contains 5 x 10 tablets in blister strips.

* Further information is available on request.



Manufactured by:
The ACME Laboratories Ltd.
Dhamrai, Dhaka, Bangladesh

For Health, Vigour and Happiness

07 1335/02