

# Laxogol

100 ml  
Oral Solution

Macrogol & Electrolytes

## Composition

Each 25 ml concentrated oral solution contains Macrogol (3350) BP 13.125 g, Sodium Bicarbonate BP 178.500 mg, Sodium Chloride BP 350.700 mg & Potassium Chloride BP 46.600 mg.

## Pharmacology

Macrogol (3350) exerts an osmotic action in the gut, which induces a laxative effect. Macrogol (3350) increases the stool volume, which triggers colon motility via neuromuscular pathways. The physiological consequence is an improved propulsive colonic transportation of the soften stools and a facilitation of the defecation. Electrolytes combined with Macrogol (3350) are exchanged across the intestinal barrier (mucosa) with serum electrolytes and excreted without net gain or loss of sodium, potassium and water. Macrogol (3350) is unchanged along the gut. It is virtually unabsorbed from the gastrointestinal tract.

## Indication

For use in adults and children (12-18) years of age for effective relief form constipation and treatment of chronic constipation. It is also effective in resolving fecal impaction, defined as refractory constipation with fecal loading of the rectum and colon.

## Dosage & Administration

Measure 25 ml of **Laxogol** oral solution with measuring cup provided, then add this to 100 ml of water (total volume 125 ml). Any unused diluted solution should be discarded within 24 hours.

### Adults :

**Constipation** : 25 ml of **Laxogol** oral solution added to 100 ml of water once daily (to make a total volume of 125 ml). This may be increased to 2-3 doses of 25 ml daily (each 25 ml dose added to 100 ml of water) if required according to individual response.

**Fecal Impaction** : 8 doses of 25 ml daily (each 25 ml dose added to 100 ml of water).

A course of treatment for fecal impaction does not normally exceed 3 days.

### Children (12 – 18 years) :

25 ml of **Laxogol** oral solution added to 100 ml of water once daily.

OR AS DIRECTED BY THE PHYSICIAN.

## Contraindications

Intestinal perforation or obstruction due to structural or functional disorder of the gut wall, ileus, severe inflammatory conditions of the intestinal tract, such as Crohn's disease and ulcerative colitis and toxic megacolon. Hypersensitivity to the active substances.

## Precautions

This medicinal product contains 8.125 mmol of sodium in each dose of 25 ml. The sodium content of **Laxogol** oral solution should be taken into consideration when administering the product to patients on a controlled sodium diet.

## Side Effects

In the treatment of chronic constipation, diarrhea or loose stools normally respond to a reduction in dose. Diarrhea, abdominal distension, anorectal discomfort and mild vomiting are more often observed during the treatment for fecal impaction. Vomiting may be resolved if the dose is reduced or delayed.

## Use in Pregnancy & Lactation

Clinically, no effects during pregnancy are anticipated, since systemic exposure to Macrogol (3350) is negligible. **Laxogol** oral solution can be used during pregnancy.

No effects on the breastfed newborn/infant are anticipated since the systemic exposure of the breast-feeding woman to Macrogol (3350) is negligible. **Laxogol** oral solution can be used during breast-feeding.

## Drug Interactions

There is a possibility that the absorption of other medicinal products could be transiently reduced during use with Macrogol & Electrolytes oral solution. There have been isolated reports of decreased efficacy with some concomitantly administered medicinal products, e.g. anti-epileptics.

## Overdose

Extensive fluid loss by diarrhea or vomiting may require correction with electrolytes.

## Storage

Store in a cool and dry place protected from light.

Keep all medicines out of reach of children.

## Supply

Each bottle contains 100 ml concentrated oral solution with a measuring cup.

\* Further information is available on request.

07 1369



Manufactured by:

**The ACME Laboratories Ltd.**

Dhamrai, Dhaka, Bangladesh

For Health, Vigour and Happiness