

Patrangasav Ayurvedic Medicine

#### Composition

Each 5 m	l contains	extracts	of
----------	------------	----------	----

Caesalpinia sappan	4.47	mg	
Acacia catechu	4.47	mg	
Adhatoda vasica	4.47	mg	
Salmalia malabarica	4.47	mg	
Sida cordifolia	4.47	mg	
Semecarpus anacardium	4.47	mg	
Hemidesmus indicus	4.47	mg	
Ichnocarpus frutescens	4.47	mg	
Hibiscus rosa sinensis	4.47	mg	
Mangnifera indica	4.47	mg	
Berberis aristata	4.47	mg	
Swertia chirata	4.47	mg	
Papaver somniferum	4.47		
Cuminum cyminum	4.47		
Aegle marmelos	4.47	mg	
Scirpus kysoor	4.47		
Cinnamomum zeylanicum	4.47		
Crocus sativus	4.47		
Syzygium aromaticum	4.47		
Roshot	4.47		
Iron dust	4.47		
Vitis vinifera	4.47	mg	
Woodfordia fructicosa	4.47	mg	
and other ingredients with excipients as required.			

# Ref:Patrangasav, BNAF-2011

## Description

Lecosav® (Patrangasav) is a poly-herbal formulation combination with time tested selected medicinal plants having proven efficacy as uterine tonic.

## Pharmacology

Acacia catechu: It contains catechin, taxifolin and catechutannic acid, catechin is biologically highly active. It is used as a hemostatic. Taxiflin has antibacterial antifungal and anti-inflammatory properties.

Adhatoda vasica: It contains vasicine, alkaloids and essential oil. It is a rich source of vitamin C and carotene, It acts as Uterine tonic.

Sida cordifolia: It contains ephedrin, phytosterols and potassium nitrate. The phytosterols have estrogenic activity. It is useful in diseases like cystitis, leucorrhoea and chronic dysentery.

Hemidesmus indicus: It has antibacterial, antiinflammatory and diuretic properties. It is used in urinary disease, leucorrhoea.

Mangnifera indica: It is astringent, anthelmintic and used in diarrhoea, uterine haemorrhage and other discharge.

Berberis aristata: It contains berberine, oxycanthine, & berbamine. It has anti-inflammatory & antiprotozoal properties. It is used in menorrhagia and leucorrhoea.

 $\textbf{\textit{Woodfordia bruticosa}}: It contains octacosanol, $\beta$-sitosterol, polystachoside, ellagic acid$ and polyphenols, allergic acid is known to have astringent and haemostatic properties and is useful in leucorrhoea and menorrhagia.

#### Indications & uses

- Leucorrhoea
- Cervicitis
- · Non-specific vaginitis
- · Pelvic inflammatory diseases

# Dosage & Administration

Route of Administration: Oral

Adult: 2-4 tea spoonful 2-3 times daily after meal

OR AS DIRECTED BY THE PHYSICIAN

## Contraindications

**During pregnancy** 

## **Warning & Precaution**

Do not use the drug after expiry date

### Side effects

Not yet known

# Use in Pregnancy & Lactation

Not recommended

### **Drug Interaction**

No such information available

# Overdose

Seek medical attention from your physician.

Store below 30° C temberature & in dry place, protected from light. Keep out of the reach of children.

# Packing

200 ml syrup in amber PET bottle with measuring cup

\* Further information is available on request

Manufactured by

The ACME Laboratories Ltd. Ayurvedic Division Dhulivita, Dhamrai, Dhaka, Bangladesh For Health, Vigour and Happiness



