Oral Emulsion MAGVET PLUS

Magnesium Hydroxide BP & Liquid Paraffin BP

Only for the use of animals

Composition

Each 5ml contains Magnesium Hydroxide BP 300 mg & Liquid Paraffin BP 1.25 ml.

Pharmacology

The osmotic force of Magnesium Hydroxide acts to draw fluids from the body and to retain those already within the lumen of the intestine, serving to distend the bowel, thus stimulating nerves within the colon wall, inducing peristalsis and resulting in evacuation of colonic contents. Magnesium Hydroxide acts as alkali to neutralize acid. Liquid paraffin acts as a stool lubricant.

Indication

Indicated for the treatment of constipation, bloat, acidosis, simple indigestion, impaction & abomasal ulceration.

Dosage & administration

Route of administration: oral

Large Animal

Upto 0-1 year: 100-200 ml. Upto 1-2 years: 200-300 ml. Over 2 years: 300-500 ml.

250 ml water to be added to the emulsion and shake well before administration.

Or, as directed by the Veterinary Physician.

Contraindication

Contraindicated in animal with dehydration and renal failure.

Warning & precaution

Some products have significant quantities of sodium and potassium and should be used cautiously in patients who should have these electrolytes in their diet.

Side effects

Common side effects: Magnesium Hydroxide in common with other magnesium salts may cause diarrhoea.

Rare side effects: Magnesium containing antacids can cause hypermagnesemia in patients with severe renal failure.

Use in pregnancy & lactation

In recommended dosage Magnesium Hydroxide do not have effect on pregnancy. However, risk & necessity should considered before administration.

Drug interaction

With medicine: Efficacy may be reduced In combination with Quinidine, Sodium polystyrene sulfonate, Iron salts, Isoniazide.

With food & others: No interaction were found with feed.

Overdose

If overdosed serious symptoms such as passing out or trouble breathing may be seen.

Withdrawal time

Milk- Milk must not be consumed up to 12 hour after administration of this drug.

Meat- Zero (0) Day

Storage

Store below 30° C and dry place, protected from light.

Keep all medicines out of reach of children.

Packing

500 ml Plastic bottle.

