WSP

SCZ Vet

Sulphaclozine Sodium Monohydrate INN

Only for the use of animals

Composition

Each 10 g powder contains Sulphaclozine Sodium (Monohydrate) INN 3 g.

Pharmacology

SCZ Vet is a broad spectrum Sulfonamide. It prevents conversion of para aminobenzoic acid (PABA) to Folic acid (Folic acid antagonist), while Folic acid is necessary for bacteria and coccidia.

Indications

SCZ Vet is used for the control & treatment of all types of Coccidiosis, Fowl Cholera & Fowl Typhoid in poultry.

Dosage & Administration

Route of Administration: For oral administration only.

Treatment

Broiler/Layer: 1.5 g - 2.0 g / liter of drinking water for 3-5 consecutive days for any age. In severe case, 2.5 g / liter of drinking water or 2.0 g to 2.5 g/ kg feed.

Control

Broiler: 1 g/L of drinking water at 3^{rd} weeks of age for 1^{st} time & 5^{th} weeks of age for 2^{nd} time. Each time the duration will be 3 consecutive days.

Layer: 1 g/L of drinking water at 3rd weeks of age for 1st time, 5th weeks of age for 2nd time and at the start of laying for 3rd time. Each time the duration will be 3 consecutive days. Or as directed by the registered Veterinary physician.

Contraindication

Hemorrhagic syndrome may observe in certain case.

Warning & Precaution

Please do not allow fresh water during the treatment period. After use close the sachet properly and use as early as possible.

Side Effects

Common: GI distress.

Rare: Sulphaclozine can precipitate in the urine, particularly when given at high dosages for prolonged periods.

Use in pregnancy & lactation or Lay

Not indicated in pregnancy & lactation or Lay.

Drug Interaction

With medicine: Not known. With food & others: Not known.

Overdose

An acute overdosed may cause GI distress (nausea, vomiting, diarrhea).

Withdrawal Time

Meat- 3 days, Eggs- 4 days.

Storage

Store below 30° C & dry place, protected from light. Keep all medicines out of reach of children.

Packing

10 x 10 g & 100 g Sachet.

