

Injection

# V-Plex Vet Plus

Vitamin and Mineral

## Composition

Each ml contains

|                                                               |           |         |
|---------------------------------------------------------------|-----------|---------|
| Thiamine HCl (Vitamin B <sub>1</sub> )                        | BP        | 75 mg   |
| Riboflavin (as Sodium Phosphate)<br>(Vitamin B <sub>2</sub> ) | BP        | 2 mg    |
| Pyridoxine HCl (Vitamin B <sub>6</sub> )                      | BP        | 5 mg    |
| Cyanocobalamin (Vitamin B <sub>12</sub> )                     | BP        | 120 mcg |
| Niacinamide                                                   | USP       | 100 mg  |
| D-Pantothenol                                                 | BP        | 20 mg   |
| Ferric Ammonium Citrate                                       | USP       | 15 mg   |
| Cobalt Gluconate                                              | Ph. Grade | 0.7 mg  |
| Copper Gluconate                                              | USP       | 0.2 mg  |

## Indication

**Ruminants:** Inappetence, metabolic disorders, poor growth, paralysis, polyencephalomalacia, blindness, nervous disorders, microcytic hypochromic anaemia, scaly dermatitis around the mouth & muzzle, loss of hair, skin lesions, leucopenia, weakness, scour, ataxia, paresis, anorexia, diarrhoea, convulsions, lacrimation & salivation.

**Horse:** Lethargy, anorexia, weight loss, ataxia, muscle tremors and convulsions.

**Dogs:** Anorexia, emesis, depression, paraparesis, torticollis, circling, convulsions, signs of "Black tongue" or "brown mouth", cheilosis, glossitis, gingivitis, bloody diarrhoea, cardiomyopathy.

## Dosage and administration

For intramuscular administration

**Cattle, Horse & Buffalo :** 5-8 ml injection for every 48 hours, 2-3 days.

**Calf, Goat & Sheep :** 2-3 ml injection for every 48 hours, 2-3 days.

**Small animal :** 0.5-1 ml injection for every 48 hours, 2-3 days.

Or As Directed by the Veterinary Physician.

## Contraindication

Contraindicated to animals hypersensitive to any of the ingredient of the product.

## Side effects

Common side effects: The product is well tolerated. There is no significant side effect in recommended dose.

Rare side effects: Not known.

## Warning & precaution

Recommended dose should not be exceeded.

## Use in pregnancy & lactation

The product may be used as per Veterinary physician's advice.

## Drug interaction

**With medicine:** Not known.

**With food & others:** Not known

## Overdose

Overdose may cause diarrhea.

## Withdrawal time

Meat & milk- 0 (zero day).

## Storage

Store below 30° C temperature and dry place, protected from light.

Keep all medicines out of reach of children.

## Packing

10 ml and 100 ml Vial.



Manufactured by

*The ACME Laboratories Ltd.*  
**Veterinary Division**

Dhaka, Bangladesh

