

WSP

Vitamix-C Vet

Vitamin C

Only for the use of animals

Composition

Each gram powder contains Ascorbic Acid (Vitamin C) 980 mg.

Indication

Poultry: For the prevention & supportive therapy of different types of stress, such as- heat stress, transportation, vaccination, hemorrhage & stress during debeaking. Helps for healing intestinal ulcer in case of coccidiosis & increase immunity.

Fish & Shrimp: For the supportive treatment of ulcerative syndrome of fish, anorexia, scoliosis & lordosis including body deformities, reducing hatching rate of brood fish. helps to increase immunity of fish & shrimp. It is also particularly effective in reducing stress during fry transportation after fry stocking.

Dosage & Administration

For oral administration only.

Poultry: 1 g/5-10 liter drinking water or 2 g/5-10 Kg feed for 5 days.

Or as directed by the registered Veterinary physician

Fish: 1-2 g/ kg feed

Or as directed by the Fisheries expert.

Warning & Precaution

Do not smell the powder. Please close the sachet after each use.

Withdrawal Time

Meat & egg- 0 (Zero) day.

Storage

Store below 30° C temperature & dry place, protected from light.

Keep out of reach of children.

Packing

100 g & 1 kg sachet



Manufactured by

The ACME Laboratories Ltd.

Veterinary Division

Dhulivita, Dhamrai, Dhaka, Bangladesh

