

W=100 mm

Imegli

Tablet

Imeglimin Hydrochloride INN

Composition

Each film coated tablet contains Imeglimin Hydrochloride INN 500 mg.

Pharmacology

Imeglimin Hydrochloride is an orally administered, first-in-class glimin. The glimins are a novel class of glucose-lowering agents that target multiple components of diabetes-associated pathology. It is an oxidative phosphorylation blocker that acts to inhibit hepatic gluconeogenesis, increase muscle glucose uptake, and restore normal insulin secretion.

Indication

Imegli (Imeglimin Hydrochloride) is used for the treatment of Type 2 Diabetes Mellitus as a monotherapy adjunct to diet and exercise.

Dosage & Administration

Route of administration: Oral.

The usual adult dosage is 500 mg of Imeglimin Hydrochloride administered orally twice daily in the morning and evening.

OR, AS DIRECTED BY REGISTERED PHYSICIAN

Contraindications

Imeglimin tablets are contraindicated in patients with:

- Hypersensitivity to the drug or any of its components
- Severe ketosis, diabetic coma or pre-coma and also for immediate remedy in type 1 diabetes of hyperglycemias with infusion and insulin

Warning & Precautions

- Dosage reduction might be required for patients with renal impairment, eGFR <15 ml/min/1.73 m²

Side Effects

The most common side effects of Imeglimin are diarrhoea, abdominal discomfort, nausea and vomiting at higher dose.

Use in Pregnancy & Lactation

Pregnancy: The safety of this product in pregnant women has not been established. Imeglimin should be used in pregnant women who may possibly be pregnant only if the expected therapeutic benefits outweigh the possible risks associated with treatment.

Lactation: Breast-feeding must be discontinued during administration of this product in lactating women (transfer to milk in animal studies has been reported).

Drug Interaction

With medicine: No drug-drug interaction has been reported yet.

With food & others: No interaction.

Overdose

In the event of an overdose, it is reasonable to employ the usual supportive measures, e.g., remove unabsorbed material from the gastrointestinal tract, employ clinical monitoring (including obtaining an electrocardiogram), and institute supportive therapy if required.

Storage

Store below 30°C temperature and dry place, protected from light.

Keep all medicines out of reach of children.

Packing

Each box contains 2x10 Tablets in Alu-Alu blister strips.

* Further information is available on request.



Manufactured by:

The ACME Laboratories Ltd.

Dhulivita, Dhamrai, Dhaka, Bangladesh

For Health, Vigour and Happiness

07 1843/01