

Dequadin VT

Dequalinium Chloride BP

Vaginal Tablet



Composition

Each vaginal tablet contains Dequalinium Chloride BP 10 mg.

Pharmacology

Dequalinium Chloride which is the active ingredient of Dequadin VT belongs to the family of anti-infective and anti-septic drugs. Dequalinium Chloride is active against number of bacteria which are causing bacterial vaginosis.


Indication

Dequadin VT is used for the treatment of bacterial vaginosis. It is inserted into the vagina for treatment at the site of infection.

Dosage and Administration

The recommended dose is 1 vaginal tablet daily for 6 days.

Usage Guideline of



Special Directions

1. Use every evening before bed. 2. You should stop the treatment during the heavy bleeding period of your menstruation and resume afterwards. 3. Do not stop treatment early, even if you no longer have any discomfort (e.g. itching, discharge, smell). A treatment of less than 6 days may result in your symptoms returning. 4. Dequadin VT contains ingredients which do not dissolve completely, and the remains of the tablet are occasionally found in the underwear. This won't affect the effectiveness of the treatment. The tablet will not stain your underwear but for your own comfort you can use a sanitary towel or panty liner. 5. If your vagina is dry, the vaginal tablet might not dissolve and be discharged as an intact tablet. As a consequence, the treatment will fail to give the optimal result. In this case the vaginal tablet can be moistened with a drop of water before insertion.

If you have the impression that the effect of Dequalinium is too strong or too weak, talk to your doctor.

Contraindication

Hypersensitivity to Dequalinium chloride or to any of the excipients of Dequalinium Chloride. Patients having ulceration of the vaginal epithelium and portion of the cervix. Those who have not yet had their first menstruation and thus did not reach sexual maturity must not use Dequadin VT.

Warning & Precautions

Intravaginal use of soaps, spermicides and vaginal douches (vaginal washes) is not recommended during treatment with Dequadin VT. Stop the treatment during the heavy bleeding period of your menstruation and resume afterwards. As with all vaginal infections, unprotected sexual intercourse during treatment with Dequadin VT is not recommended. To minimise the exposure of the newborn to Dequadin VT, vaginal tablets should not be used within 12 hours before birth.

Side Effects:

Common side effects: Like all medicines, Dequalinium can cause side effects, although not everybody gets them. Most of the side effects are mild and short lived. Sometimes the symptoms of vaginal infections (such as itching, burning and discharge) can get worse at the beginning of the treatment, before they start to get better. The treatment should be continue with the treatment, but if the complaints persist please take advise from doctor.

Rare side effects: Vaginal bleeding, Vaginal pain, Inflammation of the vagina.

Use in Pregnancy and Lactation

Based on previous experience and since Dequadin VT acts locally, no harmful effects on pregnancy and/or the unborn and breast fed infant are expected. Dequadin VT should only be used during pregnancy and during breast feeding if medically needed.

Use in Children and Adolescents

There is insufficient experience of the use of Dequadin VT in children and adolescents under 18 years of age.

Use in women above 55 years and elderly.

There is insufficient experience of the use of Dequadin VT in women above 55 years and elderly

Drug Interaction

Not known.

Over dose

No severe side effects after an overdosage of Dequalinium Chloride has been reported.

Storage

Store below 30° C temperature & dry place, protected from light.

Keep all medicines out of reach of children.

Packing

Each box contains 1x6 tablets in Alu-PVC blister pack with an Applicator.

* Further information is available on request



Manufactured by:
The ACME Laboratories Ltd.
Dhulivita, Dhamrai, Dhaka, Bangladesh.

For Health, Vigour and Happiness