

Eye drop

Cipro-D Vet

Ciprofloxacin Hydrochloride USP & Dexamethasone BP

Only for the use of animals

Composition

Each ml contains Ciprofloxacin Hydrochloride USP equivalent to Ciprofloxacin 3 mg & Dexamethasone BP 1 mg.

Pharmacology

Ciprofloxacin inhibits DNA gyrase enzymes (specially topoisomerase-2) which causes negative supercoiling of bacterial DNA. So bacterial growth is inhibited which causes death of bacteria. Dexamethasone inhibits leukocyte infiltration at the site of inflammation.

Indications

Cipro-D Vet is used against all types of bacterial diseases of eye e.g. Acute or chronic keratitis, conjunctivitis, ophthalmitis caused by gram-positive & gram-negative bacteria.

Dosage & administration

2-3 drops into conjunctival sac every four to six hours. During the initial 24 to 48 hours, the dosage may be increased to 2-3 drops every two hours.

Or as directed by the registered Veterinarian.

Contraindication

Do not administer to animals with hypersensitivity to the active ingredients of the drug.

Warning and precaution

It should not be used more than 1 month after first opening of the bottle.

Side effects

Common side effects: Eye pain, redness, itching, lacrimation, blurry vision may occur.

Rare side effects: Not known.

Use in pregnancy & lactation

It may be used in pregnancy & lactation according to benefit or risk assessment

Drug interaction

With medicine: Specific drug interaction studies have not been conducted with ophthalmic Ciprofloxacin & Dexamethasone.

With food: Not known.

Overdose

If toxicity is seen then symptomatic treatment should be applied.

Withdrawal time: Meat- 7 days.

Milk-Not known.

Storage

Store below 30° C temperature & dry place, protected from light.

Keep all medicines out of reach of children.

Packing 5 ml dropper.



Manufactured by

The ACME Laboratories Ltd.
Veterinary Division

Dhulivita, Dhamrai, Dhaka, Bangladesh



074166/01