

W= 100 mm

Lulicill

Cream

Luliconazole 1%

Composition

Each gram cream contains Luliconazole INN 10 mg.

Pharmacology

Luliconazole is a topical antifungal agent. It is a broad-spectrum antifungal agent that belongs to the azole group. It acts against fungus by inhibiting the enzyme lanosterol demethylase and blocks ergosterol synthesis which is an important constituent of fungal cell membrane. Thus it exerts strong fungicidal activity both in-vitro and in-vivo against Dermatophytes, *Candida albicans*, *Malassezia spp.*, *Trichophyton* and *Epidermophyton spp.*

Indications

Lulicill is indicated for the topical treatment of Interdigital Tinea Pedis (Athlete's Foot), Tinea Cruris and Tinea Corporis.

Dosage and Administration

Route of administration : Topical

Adult:

Interdigital tinea pedis: Lulicill cream should be applied to the affected area and approximately 1 inch of the immediate surrounding area(s) once daily for two weeks.

Tinea cruris and Tinea corporis: Lulicill cream should be applied to the affected area and approximately 1 inch of the immediate surrounding area(s) once daily for one week.

Geriatric: Same as adult.

Paediatric: Safety and effectiveness have not been established.

Or as directed by the registered physician.

Contraindications

None

Warning & Precautions

Luliconazole is recommended for topical use only. It is not intended for ophthalmic, oral or intravaginal use. If there is any unusual allergic reaction with the use of Luliconazole, then treatment should be discontinued and appropriate therapy should be instituted.

Side effects

Common side effects : Application site reactions were observed in less than 1% of subjects in clinical trial.

Rare side effects : Contact dermatitis and cellulitis.

Use in Pregnancy and Lactation

Pregnancy: Pregnancy Category C. There are no adequate and well-controlled studies in pregnant women. Luliconazole should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

Lactation: It is not known whether Luliconazole is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when Luliconazole cream is applied to women who are breastfeeding.

Use in Children & Adolescents

Safety and effectiveness have not been established for the children below 12 years.

Drug Interaction

When applied to the patients with moderate to severe Tinea Cruris, Luliconazole may inhibit the activity of CYP2C19 and CYP3A4.

Overdose

No data available.

Storage

Store below 30° C and dry place, protected from light.

Keep all medicines out of reach of children.

Packing

Each tube contains 20 gm of Lulicill cream.

* Further information is available on request.



Manufactured by:
The ACME Laboratories Ltd.
Dhulivita, Dhamrai, Dhaka, Bangladesh

For Health, Vigour and Happiness

07 1844/01