

Oral Solution

Acxin Vet

Difloxacin Hydrochloride Trihydrate BP

Only for the use of animals

Composition

Each ml contains Difloxacin (as Hydrochloride Trihydrate) BP 100 mg.

Pharmacology

Difloxacin is a fluoroquinolone and broad-spectrum bactericidal antibiotic that acts by inhibiting DNA gyrase enzyme of bacteria which prevents the DNA synthesis of bacteria & results in death of bacteria.

Indication

For the treatment of CRD, Mycoplasmosis, Fowl cholera, Colibacillosis etc. caused by susceptible microorganisms in poultry.

Dosage & Administration

Route of administration: Oral

Poultry- 1ml/Liter (10 mg/kg body weight) drinking water orally for 5 days.

Or, as directed by the registered Veterinarian.

Contraindication

Contraindicated in birds hypersensitive to Difloxacin and birds with problems of Osteoporosis or Arthropathies.

Warning and precaution

Use cautiously in animals/birds with known joint problems.

Side effects

Common- Gastrointestinal disturbance.

Rare- CNS depression, seizures, facial swelling etc. are observed in high concentration.

Use in pregnancy, lactation & lay

Not known.

Drug interaction

With medicine- Do not administer with Theophylline & products containing Aluminum, Iron, Zinc, Magnesium, or Calcium.

With food & others- Not known.

Overdose

Not known

Withdrawal time

Poultry: Meat- 01 day.

Egg- Not known

Storage

Store below 30° C temperature & dry place, protected from light. Keep all medicines out of reach of children.

Packing

100 ml & 500 ml bottle.



Manufactured by

The ACME Laboratories Ltd.

Veterinary Division

Dhulivita, Dhamrai, Dhaka, Bangladesh

