

Oral Solution
Synbiozyme Vet
Enzyme, Probiotic & Prebiotic

Only for the use of animals

Composition

Each litre contains Xylanase 250,000 IU, Protease 50,000 IU, Cellulase 40,000 IU, Phytase 40,000 IU, Amylase 30,000 IU, Pectinase 30,000 IU, β -Glucanase 20,000 IU, Mannanase 20,000 IU, Lipase 10,000 IU, *L.acidophilus* 1×10^8 cfu, *S.cerevisiae* 1×10^7 cfu & FOS (Fructooligosaccharides) 10,000 ppm.

Indications

Helps to convert feed to easily absorbable form, increase nutrition absorption, promote growth of intestinal villus, promote animal feed intake and daily weight gain, eliminate harmful intestinal pathogen & reduce poultry mortality rate.

Dosage & Administration

Route of administration: For oral administration only.
Broiler: for starter: 0.5 ml/L of drinking water, for grower & finisher: 0.25 ml/L of drinking water, Layer: 0.25 ml/L of drinking water,
Cattle: 30-50 ml/head/day, Calves: 3-5 ml/head/day,
Or, as directed by the registered Veterinarian.
Aquaculture: 250-500 ml/MT of feed.
Or, as advised by the Fisheries Expert.

Contraindication

Not known.

Warning & Precaution

Antibiotics, Acidifiers may destroy the activity of enzyme preparations.

Side Effects

Common: Not known.

Rare: Not known.

Use in Pregnancy, Lactation & Lay

Usually safe at recommended dose. However, risk & necessity should consider before administration.

Drug Interaction

With medicine: Not known.

With food & others: Not known.

Overdose

Generally safe at recommended dose.

Withdrawal Time

Meat, Milk & Egg: 0 (Zero) day.

Storage

Store below 30° C temperature and dry place, protected from light.
Keep all medicines out of reach of children.

Packing

100ml, 500 ml, & 1 L.

Marketed by

The **ACME** *Laboratories Ltd.*

Veterinary Division

Dhulivita, Dhamrai, Dhaka, Bangladesh



Manufactured by

YANTEN Biotechnology
(JiangSu) Ltd., CO. Suzhou City,
China